




BOOK OF ABSTRACTS



**32nd Marian Smoluchowski Symposium
on Statistical Physics
Information processing**



18-20 September, 2019, Kraków, Poland

32nd Marian Smoluchowski Symposium on Statistical Physics

18 - 20 SEPTEMBER, 2019, KRAKÓW, POLAND

Local Organizing Committee:

Jakub Barbasz

Karol Capała

Michał Cieśla

Bartłomiej Dybiec

Paweł F. Góra

Ewa Gudowska-Nowak

Lech Longa

Maciej Majka

Katarzyna Oleś

Wojciech Tomczyk

Conference Chair:

Ewa Gudowska-Nowak

Conference Secretary:

Ewa Witkowska

Conference webpage:

www.smoluchowski.if.uj.edu.pl

Posters (continued)

13

Kinetic equation for the pair distribution function in the Boltzmann gas

Ryszard Wojnar¹

¹ Instytut Podstawowych Problemów Techniki PAN

Corresponding Author: rwojnar@ippt.pan.pl

We report on a new kinetic equation for an auxiliary two-particle distribution function $f(\mathbf{k}_1, \mathbf{v}_1, \mathbf{k}_2, \mathbf{v}_2, t)$. Its general form reads

$$\left(\frac{\partial}{\partial t} + i \mathbf{k}_1 v_1 + i \mathbf{k}_2 v_2 \right) f(\mathbf{k}_1, \mathbf{v}_1, \mathbf{k}_2, \mathbf{v}_2, t) = \int_0^t d\tau \mathcal{G}(\mathbf{k}_1, \mathbf{k}_2, \tau) f(\mathbf{k}_1, \mathbf{v}_1, \mathbf{k}_2, \mathbf{v}_2, t - \tau) \quad (1)$$

with

$$\mathcal{G}(\mathbf{k}_1, \mathbf{k}_2, \tau) \equiv \int d\mathbf{v}^{N-2} d\mathbf{r}^N e^{-i\mathbf{k}_1 \mathbf{r}_1} e^{-i\mathbf{k}_2 \mathbf{r}_2} \mathcal{P} K_N e^{-\tau(1-\mathcal{P})K_N} e^{i\mathbf{k}_1 \mathbf{r}_1} e^{i\mathbf{k}_2 \mathbf{r}_2} \frac{f^{0N}(\mathbf{v}^N, \mathbf{r}^N)}{\varphi_M(v_1) \varphi_M(v_2)} \quad (2)$$

which is the scattering operator for our problem. We have applied to this problem the technique of projection operators. We find that $\mathcal{G}(t)$ is finite at $t = 0$ and can be readily expanded in Taylor series about $t = 0$. This is a distinct feature which is not present usually in other kinetic equations. Therefore the kinetic equation (1) with the kernel (2) is not only valid for long times but for arbitrarily short times, including $t = 0$, as well.

Next we set the distance between particles 1 and 2, $|\mathbf{r}_1 - \mathbf{r}_2| = a = \text{constant}$. This creates a dumbbell model of molecule, the diffusion of which is being studied.